

IRREDUCIBLE POLYNOMIALS IN $F_q[x]$ VERSUS $1 - qz$ IN $\mathbf{C}[z]$

BARRY BRENT

ABSTRACT. Let q be any prime power, let F_q be the field with q elements, and let N_n be the number of monic irreducible polynomials of degree n in $F_q[x]$. Let $z \in \mathbf{C}$ be close to zero. We show that $\prod_{n=1}^{\infty} (1 - z^n)^{N_n} = 1 - qz$.

1. INTRODUCTION

In this note, we prove the following

Theorem. *For q a prime power, let F_q be the field with q elements, and let N_n be the number of monic irreducible polynomials of degree n in $F_q[x]$. Let $s > q$. If $z \in \mathbf{C}$ and $|z| < \frac{1}{s}$, then*

$$\prod_{n=1}^{\infty} (1 - z^n)^{N_n} = 1 - qz.$$

2. PROOF OF THE THEOREM

Divisor sums are taken over positive divisors throughout. Let μ be the Möbius function, and let $N(a, n) = n^{-1} \sum_{d|n} \mu(n/d) a^d$. It is clear that the number $N(a, n)$ is positive when $a > 1$. We have the following

Lemma. *If $b > a > 1$, $z \in \mathbf{C}$ and $|z| < \frac{1}{b}$, then*

$$\prod_{n=1}^{\infty} (1 - z^n)^{N(a, n)} = 1 - az.$$

Proof of the lemma First we show that the infinite product is absolutely convergent, which is the same as showing that the series $S = \sum_{n=1}^{\infty} N(a, n) |\log(1 - z^n)|$ converges. By Möbius inversion,

$$(1) \quad a^n = \sum_{d|n} d N(a, d),$$

so $N(a, n) \leq a^n/n$. For small $|z|$ and large n we have the existence of a positive constant C such that $|\log(1 - z^n)| \leq C|z^n|$ because $\lim_{u \rightarrow 0} \log(1 - u)/u = 0$. Therefore the tail of the series S is dominated for some k by the series $\sum_{n=k}^{\infty} C \frac{a^n}{n} |z^n|$, which,

1991 *Mathematics Subject Classification.* 11F11.

Key words and phrases. finite fields, infinite products, irreducible polynomials.

under our hypotheses, converges by the ratio test. Hence the series S converges and the product $\prod_{n=1}^{\infty} (1 - z^n)^{N(a,n)}$ converges absolutely.

Now clearly the series $\sum_{n=1}^{\infty} N(a,n)z^n$ also converges absolutely under our hypotheses. We need these two facts because we are going to apply the following result (due essentially to Euler): an absolutely convergent infinite product $\prod_{n=1}^{\infty} (1 - z^n)^{f(n)/n}$ with the property that $\sum_{n=1}^{\infty} f(n)/n$ converges absolutely expands as a power series $1 + \sum_{n=1}^{\infty} r(n)z^n$ with (for $n = 1, 2, 3, \dots$)

$$(2) \quad nr(n) + \sum_{k=1}^n r(n-k) \sum_{d|k} f(d) = 0$$

([A], Theorem 14.8). Specializing equation (2) to the situation $f(n) = nN(a,n)$ and applying equation (1), we find that

$$(3) \quad nr(n) + \sum_{k=1}^n r(n-k)a^k = 0.$$

If we write $r(0) = 1$ and apply equation (3) twice with $n = 1$ and $n = 2$, we find that $r(1) = -a$ and $r(2) = 0$. Then it is easy to show that $r(n) = 0$ for all $n > 1$ by induction. This finishes the proof of the lemma.

If $q = p^n$, p prime, then $N(q,n) = N_n$, the number of monic irreducible polynomials in $F_q[x]$ ([G],[BeBl], or, for the case $q = p$, [IR]). So the theorem follows from the lemma.

REFERENCES

- [A] T.M. Apostol, *Introduction to Analytic Number Theory*, Springer-Verlag, New York Berlin Heidelberg Tokyo, 1976.
- [BeBl] J. Beachy and W. D. Blair, *Abstract Algebra, Third Edition*, Waveland Press, Inc., 2006.
- [G] C. F. Gauss, *Die Lehre von den Resten*, Arithmetische Untersuchungen, Chelsea, 1965.
- [IR] K. Ireland and M. Rosen, *A Classical Introduction to Modern Number Theory*, Springer-Verlag, New York Berlin Heidelberg, 1982.

4304 12TH AVENUE SOUTH, MINNEAPOLIS, MN 55407-3218
E-mail address: `barrybrent@member.ams.org`